


FORM PTO-1390 (REV. 12-97)		U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE		ATTORNEY'S DOCKET NUMBER 7024109PUR48	
TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. 371				U.S. APPLICATION NO. (If known, see 37 CFR 1.5)	
				09/180340	
INTERNATIONAL APPLICATION NO. PCT/US97/07663		INTERNATIONAL FILING DATE 05/06/1997 - May 6, 1997		PRIORITY DATE CLAIMED 05/06/1996 - May 6, 1996	
TITLE OF INVENTION STABLE RECOMBINANT YEASTS FOR FERMENTING XYLOSE TO ETHANOL					
APPLICANT(S) FOR DO/EO/US HO, Nancy W. Y. and CHEN, Zheng-Dao					
Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:					
<p>1. <input checked="" type="checkbox"/> This is a FIRST submission of items concerning a filing under 35 U.S.C. 371.</p> <p>2. <input type="checkbox"/> This is a SECOND or SUBSEQUENT submission of items concerning a filing under 35 U.S.C. 371.</p> <p>3. <input checked="" type="checkbox"/> This express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1).</p> <p>4. <input checked="" type="checkbox"/> A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date.</p> <p>5. <input checked="" type="checkbox"/> A copy of the International Application as filed (35 U.S.C. 371(c)(2))</p> <p style="margin-left: 20px;">a. <input type="checkbox"/> is transmitted herewith (required only if not transmitted by the International Bureau).</p> <p style="margin-left: 20px;">b. <input type="checkbox"/> has been transmitted by the International Bureau.</p> <p style="margin-left: 20px;">c. <input checked="" type="checkbox"/> is not required, as the application was filed in the United States Receiving Office (RO/US).</p> <p>6. <input type="checkbox"/> A translation of the International Application into English (35 U.S.C. 371(c)(2)).</p> <p>7. <input checked="" type="checkbox"/> Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3))</p> <p style="margin-left: 20px;">a. <input type="checkbox"/> are transmitted herewith (required only if not transmitted by the International Bureau).</p> <p style="margin-left: 20px;">b. <input type="checkbox"/> have been transmitted by the International Bureau.</p> <p style="margin-left: 20px;">c. <input type="checkbox"/> have not been made; however, the time limit for making such amendments has NOT expired.</p> <p style="margin-left: 20px;">d. <input checked="" type="checkbox"/> have not been made and will not be made.</p> <p>8. <input type="checkbox"/> A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371 (c)(3)).</p> <p>9. <input checked="" type="checkbox"/> An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)). (unsigned)</p> <p>10. <input type="checkbox"/> A translation of the annexes of the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).</p>					
Items 11. to 16. below concern document(s) or information included:					
<p>11. <input type="checkbox"/> An Information Disclosure Statement under 37 CFR 1.97 and 1.98.</p> <p>12. <input type="checkbox"/> An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.</p> <p>13. <input type="checkbox"/> A FIRST preliminary amendment. <input type="checkbox"/> A SECOND or SUBSEQUENT preliminary amendment.</p> <p>14. <input type="checkbox"/> A substitute specification.</p> <p>15. <input type="checkbox"/> A change of power of attorney and/or address letter.</p> <p>16. <input checked="" type="checkbox"/> Other items or information:</p> <p style="margin-left: 20px;">a. Request</p> <p style="margin-left: 20px;">b. International Publication</p> <p style="margin-left: 20px;">c. Response to Invitation to Correct Defects</p> <p style="margin-left: 20px;">d. PCT/IB/332,304,308</p> <p style="margin-left: 20px;">e. Notif of Receipt of Demand</p> <p style="margin-left: 20px;">f. Resp to International Search Report</p> <p style="margin-left: 20px;">g. Resp to Written Opinion</p> <p style="margin-left: 20px;">h. International Prelim Exam Report</p> <p style="margin-left: 20px;">i. Change in address of one of the Applicants</p>					

Express Mail label number EM577550042US

Date of Deposit: 5 NOVEMBER 1998

I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37CFR § 1.10 on the date indicated above and is addressed to the Assistant Commissioner for Patents, Washington, D.C. 20231.

Jonda C. Shelby
Signature of person mailing paper or fee

U.S. APPLICATION NO (if known, see 37 CFR 1.5)		INTERNATIONAL APPLICATION NO. PCT/US97/07663		ATTORNEY'S DOCKET NUMBER 7024109PUR48	
17. <input checked="" type="checkbox"/> The following fees are submitted: BASIC NATIONAL FEE (37 CFR 1.492 (a) (1) - (5)): Search Report has been prepared by the EPO or JPO \$930.00 International preliminary examination fee paid to USPTO (37 CFR 1.482) \$720.00 ⁵⁴⁰ No international preliminary examination fee paid to USPTO (37 CFR 1.482) but international search fee paid to USPTO (37 CFR 1.445(a)(2)) \$790.00 Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO \$1070.00 International preliminary examination fee paid to USPTO (37 CFR 1.482) and all claims satisfied provisions of PCT Article 33(2)-(4) \$98.00 ENTER APPROPRIATE BASIC FEE AMOUNT =				CALCULATIONS PTO USE ONLY	
Surcharge of \$130.00 for furnishing the oath or declaration later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(e)).					
CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE		
Total claims	35 - 20 =	15	x \$14.00	\$	210 ⁰⁰
Independent claims	7 - 3 =	4	x \$82.00/ea.	\$	256
MULTIPLE DEPENDENT CLAIM(S) (if applicable) 1			+ \$270.00	\$	210
TOTAL OF ABOVE CALCULATIONS =				\$	1,216 ⁰⁰
Reduction of 1/2 for filing by small entity, if applicable. A Small Entity Statement must also be filed (Note 37 CFR 1.9, 1.27, 1.28).				\$	
SUBTOTAL =				\$	1,216 ⁰⁰
Processing fee of \$130.00 for furnishing the English translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(f)).				\$	
TOTAL NATIONAL FEE =				\$	1,216 ⁰⁰
Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property +				\$	
TOTAL FEES ENCLOSED =				\$	1,216 ⁰⁰
				Amount to be refunded:	\$
				charged:	\$
a. <input checked="" type="checkbox"/> A check in the amount of \$ <u>1,216⁰⁰</u> to cover the above fees is enclosed. b. <input type="checkbox"/> Please charge my Deposit Account No. _____ in the amount of \$ _____ to cover the above fees. A duplicate copy of this sheet is enclosed. c. <input checked="" type="checkbox"/> The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No <u>23-3030</u> . A duplicate copy of this sheet is enclosed.					
NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137 (a) or (b)) must be filed and granted to restore the application to pending status.					
SEND ALL CORRESPONDENCE TO Kenneth A. Gandy WOODARD, EMHARDT, NAUGHTON, MORIARTY & MCNETT Bank One Center/Tower, Suite 3700 111 Monument Circle Indianapolis, Indiana 46204 US					
				SIGNATURE  Kenneth A. Gandy	
				25 NOV 96 #33,386 REGISTRATION NUMBER	
DOCUMENT PROCESSING					

SCANNED5

07 Rec'd PCT/PTO 05 NOV 1998

STABLE RECOMBINANT YEASTS
FOR FERMENTING XYLOSE TO ETHANOL

5

BACKGROUND

- The present invention relates generally to
- 10 genetically engineered microorganisms and in particular to unique methods for stably incorporating exogenous DNA into cells, including the incorporation of multiple copies of the exogenous DNA at reiterated DNA sequences in the host. In a preferred aspect, the invention
- 15 relates to yeasts capable of fermenting xylose (preferably cofermenting the same with glucose) to ethanol. More particularly, a preferred aspect of the invention relates to yeasts containing cloned genes encoding xylose reductase (XR), xylitol dehydrogenase
- 20 (XD), and xylulokinase (XK), which yeasts substantially retain their efficiency for fermenting xylose to ethanol even after culturing in non-selective medium for a large number of generations.
- 25 As further background, recent studies have proven ethanol to be an ideal liquid fuel for automobiles. It can be used directly as a neat fuel (100% ethanol) or as a blend with gasoline at various concentrations. The use of ethanol to supplement or replace gasoline can reduce
- 30 the dependency of many nations on imported foreign oil and also provide a renewable fuel for transportation. Furthermore, ethanol has proven to provide cleaner fuels that release far fewer pollutants into the environment than regular gasoline. For example, it has been
- 35 demonstrated that the use of oxygenated materials in gasoline can reduce the emission of carbon monoxide, a

- 2 -

harmful pollutant, into the air. Among the several oxygenates currently used for boosting the oxygen content of gasoline, ethanol has the highest oxygen content. The United States Environmental Protection Agency (EPA) has
5 shown that gasoline blended with 10% ethanol reduces carbon monoxide emissions by about 25% - 30%.

Up to now, the feedstock used for the production of industrial alcohol by fermentation has been sugars from
10 sugar cane or beets and starch from corn or other food crops. However, these agricultural crops are presently considered to be too expensive to be used as feedstock for the large-scale production of fuel ethanol. Plant biomass is an attractive feedstock for ethanol-fuel
15 production by fermentation because it is renewable, and available at low costs and in large amounts. The concept of using alcohol produced by microbial fermentation of sugars from agricultural biomass had its nascence at least two decades ago. The major fermentable sugars from
20 cellulosic materials are glucose and xylose, with the ratio of glucose to xylose being approximately 2 or 3 to 1. The most desirable fermentations of cellulosic materials would, of course, completely convert both glucose and xylose to ethanol. Unfortunately, even now
25 there is not a single known natural microorganism capable of fermenting both glucose and xylose effectively.

Yeasts, particularly *Saccharomyces* yeasts, have traditionally been used for fermenting glucose-based
30 feedstocks to ethanol, and they are still considered the best microorganisms for that purpose. However, these glucose-fermenting yeasts, including the *Saccharomyces*

- 3 -

yeasts, have been found to be unable to ferment xylose and also unable to use this pentose sugar for growth.

Recently, N. Ho et al. have developed recombinant yeasts, particularly recombinant *Saccharomyces* yeasts, capable of effectively fermenting xylose to ethanol (Ho and Tsao, 1995). More particularly, the preferred recombinant yeasts were capable of co-fermenting the two major sugar constituents of cellulosic biomass, glucose and xylose, to ethanol (Ho and Tsao, 1995). These recombinant yeasts were developed by the transformation of yeasts with a high-copy number plasmid containing three cloned genes, XR, XD, and XK, encoding three key enzymes for xylose metabolism (Figure 1). Figure 2 and Figure 3 demonstrate two of the prior-made recombinant *Saccharomyces* yeasts, designated 1400 (pLNH32) and 1400(pLNH33), capable of co-fermenting 8% glucose and 4% xylose present in the same medium almost completely to ethanol in two days. On the other hand, Figure 4 shows that the parent yeast fusion 1400 (D'Amore, et al., 1989 and D'Amore, et al., 1990) can only ferment glucose, but not xylose, to ethanol. 1400 (pLNH32) (in short LNH32) and 1400(pLNH33) (in short LNH33) were developed by the transformation of the *Saccharomyces* fusion 1400 (D'Amore, et al., 1989 and D'Amore, et al., 1990) with two of the high-copy-number plasmids, pLNH32 and pLNH33, shown in Figure 1. To date, there have been four such high-copy-number plasmids reported, pLNH31, pLNH32, pLNH33, and pLNH34 (Ho and Tsao, 1995). Each of these plasmids can transform fusion 1400 to recombinant yeasts to co-ferment both glucose and xylose with similar efficiencies.

- 4 -

Yeasts 1400 (pLNH32), 1400(pLNH33), and related recombinant xylose-fermenting *Saccharomyces*, with their xylose metabolizing genes cloned on a 2 μ -based stable
5 high-copy-number plasmid, are quite suitable for a batch process fermentation. However, in a continuous process fermentation, after prolonged culture in a glucose-rich medium (more than 20 generations), 1400 (pLNH32),
1400(pLNH33), and similar plasmid-mediated recombinant
10 yeasts lose their capability of fermenting xylose as shown in Figure 5 and Figure 6.

Generally, exogenous DNA or gene(s) can be cloned into yeasts by two separate ways. One way is to clone
15 the exogenous DNA or gene(s) into a plasmid vector containing a selectable genetic marker and a functional yeast DNA replication origin or ARS (autonomous replicating sequence) (Struhl et al., 1979; Stinchcomb et al., 1980; Chan and Tye, 1980) that allows the plasmid to
20 be able to replicate autonomously in its new host, followed by transformation of the desired yeast host with the plasmid containing the cloned DNA fragment or gene(s). The resulting yeast transformants are able to stably maintain the cloned gene in the presence of
25 selection pressure. However, such cloned gene(s) are unstable after prolonged culture in non-selective medium (in the absence of selection pressure).

Another way to clone the exogenous DNA or gene(s)
30 into a yeast host is to integrate the DNA or gene(s) into the yeast chromosome. In yeast, integrative transformation is almost always via homologous

- 5 -

recombination (Orr-Weaver, 1981). The simplest way to clone a desired gene into a yeast chromosome by integration is first to clone the desired gene into a plasmid which does not contain a replication of origin or ARS (autonomous replication sequences) but does contain a piece of the host DNA for targeting the integration to a specific site (Orr-Weaver, 1981). Transformation of the new yeast host with such an intact integrative vector will generate integrative transformants containing the desired gene cloned to the site next to the selected targeting yeast DNA sequences. However, the frequency of such integrative transformation is extremely low (1 to 10 transformants per μg DNA). Subsequently, it has been demonstrated that integrative vectors linearized within the DNA fragment homologous to the host chromosomal DNA can transform yeasts with much higher frequencies (100- to 1000-fold higher) (Orr-Weaver, 1981; Orr-Weaver and Szostak, 1983). It was suggested that double-stranded breaks, introduced by restriction enzyme digestion, are recombinogenic and highly interactive with homologous chromosomal DNA. This is particularly helpful for a complex plasmid, containing more than one yeast gene, so that one can direct the integration to a specific site by making a restriction enzyme cut within the corresponding region on the plasmid.

Another type of integration, also described as transplacement or gene disruption, makes use of double homologous recombination to replace yeast chromosomal DNA (Rothstein, 1981). Double homologous recombination vectors contain the exogenous DNA or gene(s) to be cloned and the selection marker, flanked by yeast DNA sequences

- 6 -

homologous to 5' and 3' regions of the segment of chromosomal DNA to be replaced. Prior to transformation, the vector is digested with restriction enzymes which liberate the transplacing fragment containing 5' and 3' ends homologous to the chromosomal DNA sequences at the desired integration sites. The latter strategy has become the method of choice for integrative transformation of yeast if a stable single-copy transformant is desired.

A number of strategies based on integration into reiterated chromosomal DNA have been used to generate stable multiple-copy integrants. For example, the delta sequence of yeast retrotransposon Ty (Sakai et al., 1990; Sakai et al., 1991), the highly conserved repeated sigma element (Kudla and Nicolas, 1992) and non-transcribed sequences of ribosomal DNA (Lopes et al., 1989; Lopes et al., 1991; Rossolini et al., 1992) have all been used as the target sites for multiple integration of exogenous gene(s) into yeast (Rothstein, 1991; Romanos et al., 1992).

Recent work reported in the literature on multiple integration of exogenous genes into the yeast chromosome has for the most part involved the use of either properly linearized non-replicative vectors or DNA fragments containing the desired gene(s) to be cloned and the genetic marker for selection, flanked with DNA sequences homologous to a region of yeast chromosomal DNA. Rarely, linearized replicative vectors and almost never intact replicative vectors, such as intact ARS vectors, were used to achieve such recombinant transformation. Thus,

- 7 -

since early work at the onset of developing yeast integrative transformation, (Szoatak and Wu (1979)), and despite the observation that DNA cloned on ARS vectors can integrate into the host chromosomes (Cregg et al., 5 1985; Kurtz et al., 1986), the use of intact ARS vectors (Struhl et al., 1979; Stinchcomb et al., 1980; Chan and Tye, 1980) for integration purposes has long since generally been abandoned. This has especially been true since the discovery that the double-stranded breaks 10 introduced by restriction enzyme digestion are recombinogenic (Orr-Weaver, 1981; Orr-Weaver and Szostak, 1983).

In light of this background, there remain needs for 15 more stable yeast which ferment xylose to ethanol, preferably xylose and glucose simultaneously to ethanol, and for facile and effective methods for making high copy number integrants. The present invention addresses these needs.

- 8 -

SUMMARY OF THE INVENTION

Accordingly, the present invention provides yeast containing multiple copies of stably cloned XR, XD, and
5 XK genes, which even upon culture in non-selective medium for multiple generations (e.g. greater than 20) retain their full capability to ferment xylose to ethanol. More preferably, the XR, XD, and XK genes are all fused to promoters not inhibited by the presence of glucose and
10 also not requiring the presence of xylose for their expression. Still more preferably, the yeast of the invention can co-ferment the two major constituents of cellulosic biomass, glucose and xylose, to ethanol.

15 Another embodiment of the present invention relates to the use of reiterated sequences, e.g. non-transcribed r-DNA sequences adjacent to the 5S DNA (Valenzuela et al., 1977), as homologous sequences for targeting high-copy-number integration of a DNA fragment containing XR,
20 XD, and XK into the yeast genome via homologous recombination. For example, a replicative plasmid vector including the DNA fragment flanked by the homologous sequences can be used to target integration of the DNA fragment. A preferred method of the invention includes
25 the steps of (a) transforming the cells with a replicative/integrative plasmid having exogenous DNA including a selection marker; and (b) repeatedly replicating the cells from step (a) to produce a number of generations of progeny cells while selecting for cells
30 which include the selection marker (e.g. by replicating on selective plates), so as to promote the retention of the replicative and integrative plasmid in subsequent

- 9 -

generations of the progeny cells and the formation of progeny cells having multiple integrated copies of the exogenous DNA. In a further step, the cells from step (b) can be replicated to produce a number of generations of progeny cells in the absence of selection for cells which include the selection marker, so as to promote the loss of the plasmid in subsequent generations of progeny cells (thus leaving an enriched population of the stable integrants).

The invention also provides an advantageous mode for selection and maintenance of the desired transformants. It is well known that in minimal medium all microorganisms require the presence of a carbon source, such as glucose or xylose, for growth. However, most microorganisms do not require the presence of a carbon source for growth in rich medium. Nevertheless, the present invention provides the use a carbon source as the selection pressure for the selection of transformants even in rich medium, such as YEP (1% yeast extract plus 2% peptone). The development of stable transformants, such as 1400(LNH-ST) (Figure 7), which are capable of effective fermentation of xylose after culturing in non-selective medium for essentially unlimited generations, has been greatly facilitated by the discovery that many yeasts, particularly *Saccharomyces* yeasts, do naturally require the presence of a carbon source, such as xylose or glucose, for growth even in rich medium, as shown in Figure 8.

In a broad aspect, the invention also provides a method for integrating multiple copies of exogenous DNA

- 10 -

into reiterated chromosomal DNA of cells. The method includes (a) transforming the cells with a replicative and integrative plasmid having exogenous DNA including a selection marker. The method also includes (b)

5 replicating the cells from step (a) to produce a number of generations of progeny cells while selecting for cells which include the selection marker, so as to promote the retention of the replicative and integrative plasmid in subsequent generations of the progeny cells and produce

10 progeny cells having multiple integrated copies of the exogenous DNA. In a specific application, such a method includes (i) transforming yeast cells with a replicative plasmid having exogenous DNA including a selection marker, the exogenous DNA being flanked on each end by a

15 DNA sequence homologous to a reiterated sequence of DNA of the host; (ii) repeatedly replicating the transformed yeast cells from step (i) to produce a number of generations of progeny cells while selecting for cells which include the selection marker, so as to promote the

20 retention of the replicative plasmid in subsequent generations of the progeny cells and result in progeny cells each containing multiple integrated copies of the exogenous DNA; and (iii) replicating the progeny cells from step (ii) to produce a number of generations of

25 progeny cells in the absence of selection for cells which include the selection marker, so as to promote the loss of the plasmid in subsequent generations of progeny cells and recover yeast cells each containing multiple copies of the exogenous DNA integrated into its chromosomal DNA.

30

In still another embodiment, the invention provides a yeast which ferments xylose to ethanol, the yeast

- 11 -

having multiple copies of exogenous DNA integrated into its chromosomal DNA. The exogenous DNA including genes encoding xylose reductase, xylitol dehydrogenase, and xylulokinase fused to non-glucose-inhibited promoters, wherein the yeast ferments glucose and xylose simultaneously to ethanol and substantially retains its capacity for fermenting xylose to ethanol for at least 20 generations even when cultured under non-selective conditions.

10

Another aspect of the invention relates to methods for fermenting xylose to ethanol, which include fermenting xylose-containing mediums with yeasts of the invention.

15

Another embodiment of the invention provides a plasmid vector for integrating an exogenous DNA sequence including a selection marker into chromosomal DNA of a target yeast cell. The inventive plasmid vector contains a functional yeast DNA replication origin and the exogenous DNA including the selection marker flanked on each end by a DNA flanking sequence which is homologous to a reiterated ribosomal DNA sequence of the target yeast cell. The plasmid further has a second selection marker in a position other than between the DNA flanking sequences.

20

25

30

A still further embodiment of the invention provides a plasmid vector for integrating an exogenous DNA sequence into a yeast to form stable integrants which ferment xylose to ethanol. The vector contains a functional yeast DNA replication origin and exogenous DNA

- 12 -

including genes encoding xylose reductase, xylitol dehydrogenase, and xylulokinase flanked on each end by a DNA flanking sequence which is homologous to a reiterated DNA sequence of the target yeast cell.

5

A still further aspect of the invention provides a method for forming cells having multiple integrated copies of an exogenous DNA fragment. This inventive method includes replicating cells having reiterated
10 genomic DNA and which contain a replicative and integrative plasmid containing the exogenous DNA to produce multiple generations of progeny cells while selecting for cells which include the selection marker, so as to promote the retention of the replicative and
15 integrative plasmid in subsequent generations of the progeny cells and produce progeny cells having multiple integrated copies of the exogenous DNA.

The invention provides yeasts containing stably
20 cloned genes enabling their use under non-selective conditions (e.g. continuous fermentations) to coferment xylose and glucose to ethanol, while not losing their capacity to ferment xylose. In addition, the invention provides methods and materials for forming stable,
25 multiple-copy integrants of yeast and other cells which are facile to perform and which can be controlled to modulate the number of copies of the integrated exogenous DNA. Additional embodiments, and features and advantages of the invention will be apparent from the following
30 description and appended claims.

- 13 -

BRIEF DESCRIPTION OF THE FIGURES

Figure 1 shows the restriction map of the plasmids pLNH31, -32, -33, and -34, and the genes cloned within.

5

Figure 2 shows that yeast transformant 1400 (pLNH32) (in short LNH32) can effectively coferment glucose and xylose. The conditions used for culturing the yeast and for fermenting the sugars are similar to those described in Example 7.

10

Figure 3 shows that yeast transformant 1400 (pLNH33) (in short LNH33) can effectively coferment glucose and xylose. The conditions used for culturing the yeast and for fermenting the sugars are similar to those described in Example 7.

15

Figure 4 shows that the parent yeast fusion strain 1400 can ferment glucose but not xylose. The conditions used for culturing the yeast and for fermenting the sugars are similar to those described in Example 7.

20

Figure 5 demonstrates that yeast transformant 1400 (pLNH32) (in short LNH32) with its xylose metabolizing genes cloned in the replicative plasmid pLNH 32 is not stable in a non-selective medium. After being cultured for 20 generations in a non-selective (for example, glucose) medium, 1400 (pLNH32) lost its capability to ferment xylose.

25
30

- 14 -

Figure 6 demonstrates that yeast transformant 1400(pLNH33) (in short LNH33) with its xylose metabolizing genes cloned in the replicative plasmid pLNH 33 is not stable in a non-selective medium. After
5 being cultured for 20 generations in a non-selective medium (for example, glucose medium), 1400(pLNH33) lost its capability to ferment xylose.

Figure 7 shows that yeast transformant 1400(LNH-ST)
10 (in short LNH-ST) can stably maintain its xylose fermenting capability even after being cultured in non-selective medium for more than 40 generations.

Figure 8 demonstrates that *S. cerevisiae* and other
15 *Saccharomyces* yeasts require a carbon source for growth even when rich media such as yeast extract and pepton were present in the medium. For example, these experiments showed that *S. cerevisiae* was unable to grow in the YEP medium containing 1% yeast extract and 2
20 % pepton, but was able to grow when glucose or xylulose was added to the YEP medium.

Figure 9A shows the restriction map of pLNH-ST, and the genes cloned within.
25

Figure 9B shows the genetic map (the order and orientation) of genes (5S rDNA, KK, AR, and KD) cloned in pLNH-ST. The oligonucleotides (for example, Oligo 25, Oligo 26, etc.) that are above or below the gene map are
30 the primers used to characterize the order and orientation of the cloned genes by PCR.

- 15 -

Figure 10 is a schematic diagram outlining the construction of pBluescript II KS(-) containing the cloned XR, XD, XK genes: four such plasmids were constructed. The KK-AR-KD fragment cloned in pKS(-)-KK-AR-KD-3 was chosen to be cloned in pUCKm-rDNA(5S)-ARS for the construction of pUCKm-rDNA(5S)(KRD)-ARS, also designated as pLNH-ST.

Figure 11 shows that yeast transformant 1400(LNH-ST) (in short LNH-ST), superior to 1400(pLNH 32) and 1400(pLNH 33), can effectively coferment glucose and xylose. The conditions used for culturing the yeast and for fermenting the sugars are similar to those described in Example 7.

Figure 12 shows the genes cloned in and the restriction map of a broad-host plasmid for the isolation of ARS containing DNA fragments from the chromosome DNA of *S. cerevisiae* and other yeasts.

- 16 -

DETAILED DESCRIPTION OF THE INVENTION

For the purpose of promoting an understanding of the principles of the invention, reference will now be made to certain preferred embodiments thereof, and specific language will be used to describe the same. It will nevertheless be understood that no limitation of the scope of the invention is thereby intended, such alterations, further modifications and applications of the principles of the invention as described herein are being contemplated as would normally occur to one skilled in the art to which the invention relates.

As mentioned above, one preferred aspect of the present invention provides recombinant yeasts incorporating stably cloned XR, XD and XK genes, which represent an improvement upon prior-reported recombinant yeasts. Generally, recombinant yeasts that can effectively coferment both glucose and xylose present in the same medium have been reported (Ho and Tsao, 1995). The yeasts made in this publication were accomplished by cloning properly modified XR, XD, and XK genes on a high copy number plasmid, pUCKm10, followed by using the resulting plasmid, pLNH3X (X=1 to 4) (Figure 1), to transform suitable natural yeasts. For example, the plasmids pLNH32 and pLNH33 were used to transform fusion yeast 1400 to 1400 (pLNH32) and 1400 (pLNH33), respectively. These recombinant *Saccharomyces* yeasts can effectively coferment both glucose and xylose present in the same medium to ethanol as shown in Figures 2 and 3, while the parent unengineered 1400 yeast can only ferment

- 17 -

glucose alone, not coferment both glucose and xylose (Figure 4).

Plasmid-mediated recombinant yeasts can maintain the
5 cloned genes in the presence of selection pressure, but
not in the absence of selection pressure. As
demonstrated in Figures 5 and 6, 1400 (pLNH32) and 1400
(pLNH33) eventually lose their plasmids and their
capability for fermenting xylose after prolonged culture
10 in the absence of selection pressure.

It is highly desirable that recombinant industrial
yeasts, particularly those strains used for the
production of large volume industrial products, such as
15 ethanol, be stable without requiring the presence of
selection. The development of recombinant yeasts
containing integrated XR, XD, and XK genes, as in the
present invention, provides such stability. In addition,
for the resulting recombinant yeasts to have the ability
20 to coferment glucose and xylose at efficiencies similar
to or better than 1400(pLNH32) and 1400(pLNH33), the
recombinant yeasts must contain not only the integrated
xylose metabolism genes, but also high numbers of copies
of such integrated genes. In preferred aspects of the
25 present invention, high-copy-number (hcn) integrants of
yeasts (i.e. yeasts having at least about 10 integrated
copies of the exogenous DNA) have been developed by
targeting a non-coding region, such as a non-coding
region of 5S ribosomal DNA (rDNA) as the site for
30 multiple integration.

- 18 -

rDNA provides an advantageous location for integration because it is highly conserved, and yeasts generally contain more than 100 copies of the rDNA repeated sequences. It will be understood, however, that to achieve yeasts of the present invention, it will not be necessary to achieve integration of the desired genes at every occurrence of a repeated or reiterated sequence. It will be sufficient to achieve such integration at each of multiple sites of a reiterated sequence, i.e. two or more sites, in accordance with the broad aspects of the present invention.

In order to integrate hcn XR, XD, and XK into the yeast chromosome at the site of 5S rDNA, the integration plasmid, pLNH-ST, as shown in Figure 9, was constructed. pLNH-ST is a yeast-*E. coli* shuttle vector and a derivative of pUCKm 6 plasmid (Ho et al., 1984). The 5S rDNA sequences was inserted at the Xho I restriction site of pUCKm 6. The 5S rDNA sequence was copied from the yeast chromosomal DNA by the PCR technique and modified by the site-specific mutagenesis technique to add an XhoI restriction site in its center (approximately) sequence as shown in Figure 9. The XhoI fragment from pKS(-)-KK-AR-KD (Figure 10) (Ho and Tsao, 1995) has been inserted into the XhoI site of the 5S rDNA cloned in pLNH-ST.

pLNH-ST differs from other traditional 5S rDNA-based hcn yeast integrating vectors in that it also contains a functional yeast ARS sequence (Struhl et al., 1979; Stinchcomb et al., 1980; Chan and Tye, 1980) as shown in Figure 9. Thus, pLNH-ST is both a replicative vector and an integrative vector. Uniquely, pLNH-ST functions first

- 19 -

as a replicative vector then as an integrative vector in the development of recombinant yeasts containing high copies of integrated XR, XD, and XK. The ARS fragment was inserted at the EcoR1 site of pUCKm 6. In addition, 5 pLNH-ST also contains the kanamycin resistance gene (Km^R) and the ampicillin resistance gene (Ap^R). Km^R functions as a geneticin resistance gene in yeasts and will confer its yeast transformants resistant to geneticin. The XhoI site of Km^R was removed by PCR technique without 10 affecting its activity. Both Km^R and Ap^R are part of the original pUCKm 6 plasmid.

As noted above, the above-described vectors differ from those used in state-of-the-art techniques by 15 containing an ARS sequence. In addition, in prior-reported methods for making hcn yeast integrants, integration of the cloned genes has taken place instantly, at the moment when the yeast cells are transformed with the exogenous genes. To the contrary, 20 in accordance with preferred modes of the invention, integration of the cloned genes continues to take place gradually, long after transformation has been completed. In particular, transformation is established first via the presence of replicative plasmid, such as pLNH-ST, in 25 the transformed yeast cells, and integration takes place only gradually via repeated replication of the transformants on plates containing selective medium.

Thus, this invention relates the use of the 30 following procedures to develop yeast or other cell transformants containing hcn integrated cloned gene(s).

- 20 -

Host cells which contain reiterated DNA sequences, for example yeast or eukaryotic cells, are transformed with a replicative/integrative plasmid, such as pLNH-ST, and transformants containing high-copy numbers of the

- 5 replicative/integrative plasmid are selected. The resulting selected transformants are repeatedly replicated onto fresh selective plates and grown to high cell density for a sufficient number of times to integrate the desired number of copies of the exogenous
- 10 DNA, followed by culturing the transformants in non-selective medium for a sufficient number of generations to remove the replicative/integrative plasmids from the transformants. The resulting transformants can then be cultured in selective medium, and those transformants
- 15 retaining their capability to effectively grow in selective medium will be those that contain hcn of the desired exogenous genes integrated into the chromosome of the yeast or other host cells. For example, fusion 1400 yeast has been transformed with pLNH-ST according to the
- 20 above described procedures, and the resulting stable recombinant yeast, 1400(LNH-ST), can coferment both glucose and xylose better than 1400(pLNH 32) and 1400(pLNH33), as shown in Figure 11. Importantly, the newly-developed stable recombinant yeast, 1400(LNH-ST),
- 25 can still ferment both glucose and xylose with equal efficiencies after being cultured in non-selective medium for 4, 20, and 40 generations as shown in Figure 7, while 1400(pLNH 32) and 1400(pLNH33) will lose most of their activity for fermenting xylose after 20 generations of
- 30 being cultured in non-selective medium (Figures 5 and 6). Furthermore, 1400(LNH-ST) has subsequently been cultured in non-selective medium for several hundred generations,

- 21 -

and still retains its full activity in cofermenting both glucose and xylose.

In the preferred methods for developing stable hcn
5 integrants, a common selection marker(s) is/are used for
the selection and maintenance of both the plasmid-
mediated activity and the activity contributed by the
integrated genes with the same selective medium. In the
present work, the common selection markers are the three
10 cloned xylose metabolism genes, XR, XD, and XK, and the
common selective medium is either rich or minimal medium
(for yeasts) containing xylose. In addition, these
cloned genes serve as the selection markers in rich
medium for most *Saccharomyces* yeasts, since the
15 applicants have shown that most of the *Saccharomyces*
yeasts do require the presence of a carbon source, such
as xylose, for growth even in rich medium (Figure 8).
Although it is not crucial for yeasts chosen as hosts to
require the presence of a carbon source in rich medium
20 for growth, it is, nevertheless, much more convenient to
be able to select the desired integrants on plates
containing rich medium with xylose rather than on plates
containing minimal medium with xylose. Preferred hosts
for transformation in the present invention belong to the
25 *Saccharomyces* species, since they are usually
extraordinarily effective for fermenting glucose. In the
event that species of yeasts desired for use as hosts for
integrating high copy numbers of xylose metabolizing
genes are found not to require the presence of a carbon
30 source for growth in rich medium, a suitable mutant of
that species which does require the carbon source in rich
medium can be isolated using conventional procedures.

- 22 -

The replicative/integrative plasmid, such as pLNH-ST, for achieving hcn integration also desirably contains a second selection means for the selection of replicative plasmid-mediated transformants. For pLNH-ST, the second selection mechanism utilizes both Km^R and Ap^R as selectable markers. Although it is not crucial for a replicative/integrative vector to contain a second selection system, it will provide more preferred vectors, particularly if the ARS vector is not sufficiently stable even in the presence of the selection pressure, and the transformants have the tendency to lose most of their plasmids prior to integrating sufficient copies of the desired genes. When using vectors which contain a second selection mechanism, the transformants may be cultured in the presence of the second selective reagent to boost their plasmids' copy number, or to re-transform the transformants with the same vector but using the second selection mechanism to re-select the transformants so that the integration process can be continued or re-initiated.

The use of both Km^R and Ap^R as the second selection system is desirable for the applicants' preferred yeast.

Km^R can be a dominant selection marker for transforming yeasts that are resistant to geneticin, but some yeasts are naturally resistant to geneticin without acquiring the plasmid containing Km^R . As a result, Km^R alone is not a preferred selection marker for the selection of yeast transformants. On the other hand, Ap^R can be effectively expressed in most yeasts, but it generally

- 23 -

cannot be used as a dominant selection marker for yeast transformation because most yeasts are naturally resistant to ampicillin. However, both Km^R and Ap^R together serve as an excellent dominant selection system for most yeasts, particularly the *Saccharomyces* yeasts. To use such a selection system, the transformants are first selected on plates containing YEPD (1% yeast extract, 2% peptone, 2% glucose) and proper concentrations of geneticin (20-80 μ g/ml, varying from species to species). The resulting transformants are screened for the expression of the Ap^R by the penicillinase test (Chevallier and Aigle, 1979) to identify true transformants.

The presence of Ap^R in pLNH-ST (Figure 9) and related replicative/integrative plasmids also serves another function. Since Ap^R is only present in the replicative plasmid and not present on the fragment integrated into the yeast chromosome, the ampicillin test also serves as a convenient process for identifying those transformants containing hcn integrated cloned genes but not plasmid vectors.

A feature of the inventive approach for providing stable recombinant yeasts containing hcn integrated gene(s) is that the number of copies of the gene(s) to be integrated can easily be controlled. For example, more copies of the XR-XD-XK genes can be inserted into the fusion yeast 1400 chromosome if another selection marker, such as Km^R , is inserted into the 5S rDNA fragment (or the targeting sequence). Furthermore, the inventive

- 24 -

methods for the development of hcn yeast integrants are also easier to accomplish than other reported approaches, wherein experimental conditions may have to be adjusted and controlled and the transformation process may have to
5 be repeated before a stable strain could be obtained.

Thus, the applicants have improved upon the stability of prior recombinant xylose-fermenting yeasts, such as 1400 (pLNH32) and 1400 (pLNH33), and developed
10 advantageously stable recombinant yeasts, for example 1400(LNH-ST), that will not require the presence of selection pressure to maintain the cloned genes and are also as effective as or even more effective for cofermenting glucose and xylose than 1400(pLNH32) and
15 1400(pLNH33). Furthermore, the applicants have also developed a convenient method that has provided the facile hcn integration of exogenous gene(s) into the cellular chromosome, wherein the number of copies of the gene(s) to be integrated is also readily controllable.

20 Similar to 1400 (pLNH32) and 1400(pLNH33), the preferred stable genetically engineered xylose-fermenting yeasts of the invention can also effectively coferment both glucose and xylose. This is because the XR, XD, and
25 XK genes inserted into the chromosome of the new yeast hosts are all fused to intact 5' non-coding sequences from genes that can be efficiently expressed in yeast, encoding the production of high levels of enzymes, and also which are not inhibited by the presence of glucose
30 in the medium. For example, the intact 5' non-coding DNA sequences that contain all the genetic elements for efficient expression of the glycolytic genes and for the

- 25 -

production of high levels of glycolytic enzymes are suitable as replacements for the intact 5' non-coding sequences of XR, XD, and XK for these purposes.

- 5 The XR, XD, and XK cloned on pLNH-ST are from *Pichia stipitis* (XR and XD) and *Saccharomyces cerevisiae*. (XK). However they can be from any microorganisms as long as they can produce high levels of the respective enzymes after they have been fused to
- 10 the proper 5' non-coding sequences containing effective promoters, ribosomal binding sites, etc. For example, these three genes are well known to occur in a wide variety of microorganisms and numerous XR, XD and XK genes have been identified and isolated. The particular
- 15 source of these genes is thus not critical to the broad aspects of this invention; rather, any DNAs encoding proteins (enzymes) having xylose reductase activity (the ability to convert D-xylose to xylitol with NADPH and/or NADH as cofactor), xylitol dehydrogenase activity (the
- 20 ability to convert xylitol to D-xylulose with NAD⁺ and/or NADP⁺ as cofactor), or xylulokinase activity (the ability to convert D-xylulose to D-xylulose-5-phosphate) will be suitable. These genes may be obtained as naturally-occurring genes, or may be modified, for
- 25 example, by the addition, substitution or deletion of bases to or of the naturally-occurring gene, so long as the encoded protein still has XR, XD or XK activity. Similarly, the genes or portions thereof may be synthetically produced by known techniques, again so
- 30 long as the resulting DNA encodes a protein exhibiting the desired XR, XD or XK activity.

- 26 -

As examples, suitable sources of XR and XD genes include xylose-utilizing yeasts such as *Candida shehatae*, *Pichia stipitis*, *Pachysolen tannophilus*,

5 suitable sources of XK genes include the above-noted xylose-utilizing yeasts, as well a xylose non-utilizing yeasts such as those from the genus *Saccharomyces*, e.g. *S. cerevisiae*, the genus *Schizosaccharomyces*, e.g. *Schizosaccharomyces pombe*, and bacteria such as
10 *Escherichia coli*, *Bacillus* species, *Streptomyces* species, etc. Genes of interest can be recovered from these sources utilizing conventional methodologies. For example, hybridization, complementation or PCR techniques can be employed for this purpose.

15 A wide variety of promoters will be suitable for use in the invention. Broadly speaking, yeast-compatible promoters capable of controlling transcription of the XR, XD or XK genes will be used.

20 Such promoters are available from numerous known sources, including yeasts, bacteria, and other cell sources. Preferably, the promoters used in the invention will be efficient, non-glucose-inhibited promoters, which do not require xylose for induction.

25 In this regard, an "efficient" promoter as used herein refers to a 5' flanking sequence which provides a high level of expression of the fused gene. Promoters having these characteristics are also widely available, and their use in the present invention, given the teachings
30 herein, will be within the purview of the ordinarily skilled artisan, as will be the fusion of the promoters to the XR, XD and XK genes, the cloning of the

- 27 -

promoter/gene fusion products into appropriate vectors and the use of the vectors to transform yeast. All of these manipulations can be performed using conventional genetic engineering techniques well known to the art and
5 literature.

The yeast DNA replication origin, e.g. the ARS containing DNA fragment, can be obtained from yeast chromosomal DNA or from chromosomal DNA of other
10 organisms, so long as the DNA fragment can function as an active replication origin to support autonomous replication of plasmid in the host chosen for hcn integration. DNA fragments which function as ARSs can readily be isolated by incorporating randomly-digested
15 DNA fragments into an *E. coli* plasmid, followed by transformation of the desired host organism, e.g. a *Saccharomyces* yeast, with the resulting bank of plasmids, as reported in the literature (Stinchcomb et al., 1980; Ho et al., 1984).

20 Novel methods have been used to create the stable strains of the present invention. Nevertheless, there are several lines of evidence indicating that the cloned genes are not on a replicative plasmid and have been
25 integrated into the host genome. For example, chromosomal DNA isolated from 1400(LNH-ST) can be used as template for the isolation of the cloned genes, including the fusions containing both the 5s rDNA and the cloned gene sequences, by the polymerase chain reaction (PCR).
30 Also, while few plasmids (pLNH-ST) can be recovered from 1400(LNH-ST) via transformation of *E. coli* (Ward, 1990), under the same conditions, hundreds of pLNH32 or pLNH33

- 28 -

plasmids can be recovered from 1400 (pLNH32) and 1400(pLNH33), respectively. Furthermore, the initial 1400 fusion yeast transformants containing high copy numbers of the replicative plasmid pLNH-ST are unstable (with respect to their capability to ferment xylose) but positive for penicillinase (enzyme encoded by Ap^R) test (Chevallier and Aigle, 1979). On the contrary, the final stable transformants, 1400(LNH-ST), which retain their capability for fermenting xylose without the presence of selection, are found to be negative for penicillinase test. This is expected if the exogenous DNA is integrated at the site of 5S rDNA since Ap^R is not part of the DNA fragment to be integrated into the host chromosome. It is also possible that some of the stable yeast transformants may contain exogenous genes integrated at the ARS sites of the yeast chromosome.

For purposes of promoting a further understanding of the present invention and its features and advantages, the following Examples are provided. It will be understood, however, that these Examples are illustrative, and not limiting, of the invention.

EXAMPLE 1

SYNTHESIS OF THE 5S rDNA FRAGMENT BY PCR

For the synthesis of the 5S rDNA fragment by PCR (to serve as the yeast DNA sequence for targeting high-copy-number integration into the yeast chromosome), the following oligonucleotides were synthesized and used as the primers for PCR reactions according to the published 5S rDNA sequence (Valenzuela et al., 1977). In addition

- 29 -

to the 5S rDNA sequence, additional nucleotides specifying the Sal I restriction site were also added to the 5' terminal of primers to facilitate the cloning of the PCR synthesized 5S rDNA into an *E. coli* plasmid.

5

Oligonucleotide I: TTAGTCGACGTCCCTCCAAATGTAAAATGG.

Oligonocleotide II: AATGTCGACGTAGAAGAGAGGGAAATGGAG

10

Chromosomal DNA isolated from fusion yeast 1400 was used as the template for the PCR reaction. The PCR synthesized 5S rDNA fragment was first cloned into the *E.coli* pBluescript II KS(-) plasmid (Stratagene Cloning Systems, La Jolla, CA) at its SalI site. The resulting

15

plasmid was designated as pKS-rDNA(5S).

EXAMPLE 2

INSERTION OF XHOI SITE INTO CLONED 5S rDNA SEQUENCE

20

The nucleotide sequence between -29 and -56 of the 5S rDNA sequence (Valenzuela et al., 1977) was modified by oligonucleotide-mediated site-specific mutagenesis (Kunkel, 1985; Kunkel et al., 1987). As a result, an XhoI restriction site was inserted at the specific site

25

described above. The protocol provided by Bio-Rad Laboratories, Inc. for oligonucleotide-mediated site-specific mutagenesis was followed to accomplish this task, except that pKS plasmid was used rather than plasmid pTZ18U or pTZ19U. The resulting plasmid

30

containing the mutated 5S rDNA was designated as pKS-5S rDNA(XhoI). The following oligonucleotide was used to

- 30 -

carry out the site-specific mutagenesis:
GAGGGCAGGCTCGAGACATGTTTCAGTAGG.

EXAMPLE 3

5

ISOLATION OF DNA FRAGMENTS FROM *S. CEREVISIAE* DNA OR OTHER DNA FUNCTIONING AS ARS IN YEASTS

10

S. cerevisiae DNA (or DNA from other yeasts or other organisms) was digested with Sau3A restriction enzyme and cloned into the Bam H1 site of pUCKm6 (Figure 12) (Ho, et al., 1984). The resulting bank of plasmids was used to transform *S. cerevisiae*. Those transformants that were capable of growing on plates containing YEPD (1% yeast extract, 2% peptone, and 2% glucose) and 50 µg/ml

15 geneticin and which were also positive for the penicillinase test (Chevallier and Algle, 1979) were selected. The plasmids from the selected true transformants were recovered by a procedure similar to that described by Ward (1990).

20

The yeast DNA fragments inserted in pUCKm6 (Figure 12) and recovered from the yeast transformants should all contain a segment of DNA that can function as an ARS (autonomous replicating sequence) in *S. cerevisiae*,

25 possibly in other yeasts as well. The DNA inserts were digested with various restriction enzymes and the resulting DNA fragments were re-inserted into pUCKm6. The latter plasmids were used to retransform *S. cerevisiae*. Any properly-sized restriction fragments

30 that can make pUCKm6 function effectively as a yeast plasmid must contain an effective "ARS" and can be used

- 31 -

to construct replicative/integrative vectors such as pLNH-ST for high-copy-number integration of exogenous gene(s) into the chromosomes of *S. cerevisiae*. These restriction fragments are also likely to function as
5 ARS's in other yeasts, and are suitable for the construction of replicative/integrative plasmids for other yeasts.

EXAMPLE 4

10 REMOVAL OF THE XhoI RESTRICTON SITE FROM THE GENETICIN (KANAMYCIN) RESISTANCE GENE, Km^R

The geneticin (kanamycin) resistance gene, Km^R , from Tn 903 (A. Oka et al., 1981) and the 5S rDNA fragment
15 described in Example 1 are part of the plasmid designed for the integration of multiple copies of exogenous genes into the yeast chromosome. However, Km^R contains an XhoI site in its coding sequence. This is in conflict with the fact that an XhoI site has been engineered into the
20 center of the cloned 5S rDNA sequence to be used for inserting exogenous genes such as XR, XD, and XK into the plasmid for integration. Thus, it is necessary to remove the XhoI site from Km^R . This can be accomplished by a number of different approaches. The applicants chose to
25 use site-specific mutagenesis by the overlap extension PCR technique (S. N. Ho, et al., 1989) to remove the XhoI site from Km^R without changing its amino acid coding sequence and without affecting the catalytic activity of the enzyme encoded by the gene. The Km^R gene cloned in
30 pUCKm6 (Figure 12) was converted to $Km^R(-Xho)$ as described above.

- 32 -

The four oligonucleotides used to accomplish this task are listed below.

Oligonucleotide I: GGCCAGTGAATTCTCGAGCAGTTGGTG

5

Oligonucleotide II: TGGAATTTAATCGCGGCCCCCTAGCAAGACG

Oligonucleotide III: TTACGCCAAGCTTGGCTGC

10 Oligonucleotide IV: TTCAACGGGAAACGTCTTGCTAGGGGCCGC

pUCKm6 (Figure 12) is a derivative of pUC9. Part of Oligo I and the entire Oligo III are synthesized according to the sequence of the polylinker region of pUC9 (Sambrook, et al. 1989).

15

The above-described genetic manipulation of pUCKm6 not only resulted in the deletion of the XhoI restriction site from the coding region of Km^R but also inserted an XhoI restriction site between the Km^R coding sequence and the EcoRI site of pUCKm6. The resulting plasmid was designated as pUCKm(-XhoI)(+XhoI). The addition of an XhoI site downstream to the Km^R coding sequence is to facilitate the insertion of the 5S rDNA fragment described in Example 1 into the newly developed plasmid pUCKm(-Xho)(+Xho).

20

25

EXAMPLE 5

CONSTRUCTION OF PLASMID pLNH-ST

30

The plasmid pUCKm(-XhoI)(+XhoI) described in Example 4 was used for the construction of pLNH-ST, shown in

- 33 -

Figure 9. First, the Sal I fragment containing the 5S rDNA(XhoI) was isolated from pKS-5S rDNA(XhoI) and inserted at the XhoI site of pUCKm(-XhoI)(+XhoI). The resulting plasmid was designated as pUCKm-rDNA(5S). To the latter plasmid, an EcoRI fragment containing an effective ARS isolated from *S. cerevisiae* (according to the procedure described in Example 3) was inserted into the EcoRI site of pUCKm-5S rDNA, and the resulting plasmid was designated as pUCKm-5S rDNA-ARS. To the latter plasmid, the XhoI fragment from pKS(-)-KK-AR-KD-3 containing the cloned XR, XD, and XK fused to yeast alcohol dehydrogenase promoter (XR), and pyruvate kinase promoter (for both XD & XK), were inserted into the XhoI site located at the center of the cloned 5S rDNA sequence. The resulting plasmid, pUCKm-rDNA(5S)(KDR)-ARS, also designated pLNH-ST, shown in Figure 9.

EXAMPLE 6

20 TRANSFORMATION OF FUSION YEAST 1400 WITH PLNH-ST AND SELECTION OF STABLE TRANSFORMANTS 1400 (LNH-ST)

pLNH-ST was used to transform fusion strain 1400 by electroporation under the conditions used for transformation of strain 1400 by plasmids pLNH32 and pLNH33 (International Publication No. 95/13362, May 18, 1995, publishing International Application No. PCT/US94/12861, filed November 8, 1994). Briefly, fifty ml yeast cells, grown to early log phase (Klett Unit (KU) 140-190), were centrifuged to remove the medium, washed twice with cold water, once with cold 1 M sorbitol, and resuspended in 200 µl 1 M sorbitol. Sixty µl of the cells were transferred into a 4 ml presterilized plastic

- 34 -

tube (with cap) and to which 1 µg plasmid DNA was added. Fifty µl of the resulting cells and plasmid mixture were pipetted into a precooled gene pulser cuvette with a 0.2 cm electrode gap and the content in the cuvette was
5 subjected to pulse by the gene pulser with a pulse controller (BioRad) at 2.0 KV, 25 µF, 200 ohms.

Immediately, .50 ml YEPD was added to the cuvette. The content of the cuvette was transferred to a new 4 ml
10 sterilized plastic tube and incubated at 30°C for 1 hr. 100 µl of the cells were plated on agar plates containing YEPD and 40 µg/ml G418 (geneticin). Fast growing colonies were selected and replicated on another plate containing the same medium. The selected colonies were
15 subjected to the ampicillin test (Chevallier and Aigle, 1979) until a positive one was identified. The above-described electroporation procedure is based on that reported by Becker and Guarente (1971).

20 Once a transformant had been positively identified by the penicillinase test, it was maintained on a YEPX (1% yeast extract, 2% peptone, 2% xylose) plate. Initially, the transformants were very unstable. They lost their xylose fermenting capability if cultured in
25 YEPD medium over 20 generations. However, by continuing to culture the transformants to stationary phase on YEPX plates, and repeatedly transferring them to fresh YEPX plates, the transformants gradually became stable with regard to their capability to ferment xylose. Once
30 stable, the transformants could be cultured in non-selective medium for several hundred or more generations

- 35 -

and were still capable of co-fermenting both glucose and xylose, as demonstrated in Example 8.

EXAMPLE 7

5 Co-FERMENTATION OF GLUCOSE AND XYLOSE WITH 1400 (LNH-ST)

A mixture of glucose and xylose (approximately 10% glucose and 5% xylose) was fermented by strain 1400 (LNH-ST) under the conditions described below. The seed
10 cultures of 1400 and 1400 (LNH-ST) were cultured aerobically in liquid YEPD medium until mid-log phase (between 400-450 Klett Units (KU)) and stored at 4°C. New seed cultures were prepared once a month by transferring 2 ml of the culture to 50 ml of fresh YEPD
15 and cultured as described above. 2 ml of the seed cultures of 1400 (LNH-ST) were inoculated into 100 ml of YEPD medium in a 300 ml Erlenmeyer flask equipped with a side-arm which allowed direct monitoring of the growth of the yeast cultures by the Klett colorimeter. The culture
20 was incubated in a shaker at 30°C and 200 rpm aerobically.

When the cell density reached mid-log phase (400-450 KU), 20 ml (50%) glucose and 10 ml (50%) xylose were
25 added to the flask. After thorough mixing, 1 ml of the culture mixture was removed from the flask to serve as the zero sample. The flask was then sealed with Saran wrap to allow fermentation to be carried out anaerobically. One ml samples of the fermentation broth
30 were removed at proper intervals (every 6 hrs.) to serve as samples for measuring glucose, xylose, xylitol, and glycerol contents of the broth during fermentation by

- 36 -

HPLC as described in Example 9. The results, shown in Figure 11, demonstrate that the genetically engineered yeast 1400 (LNH-ST) can co-ferment most of the 10% glucose and 5% xylose to ethanol in 30 hrs. The fermentation was carried out under normal conditions, without requiring special medium or pH, and also without requiring growth of yeast to high cell density. Thus, the genetically engineered 1400 (LNH-ST) can effectively co-ferment high concentrations of both glucose and xylose to ethanol with very little xylitol produced as a by-product. In comparison to the recombinant *Saccharomyces* 1400 (pLNH32) and 1400 (pLNH33) shown in Figures 2 and 3, 1400 (LNH-ST) co-fermented both glucose and xylose somewhat better than the two previously developed yeasts.

EXAMPLE 8

COMPARISON OF THE STABLE STRAIN 1400 (LNH-ST) WITH 1400 (LNH32) AND 1400 (LNH33) IN CO-FERMENTING GLUCOSE AND XYLOSE AFTER CULTURE IN NON-SELECTIVE MEDIUM FOR 4, 20, AND 40 GENERATIONS.

As described in Example 7, 2 ml each of the seed cultures of 1400 (LNH-ST), 1400 (LNH32), and 1400 (LNH33) were inoculated into 50 ml YEPD in separate 250 ml Erlenmeyer flasks equipped with side-arms. After the cells were cultured to 400-450 KU, 2 ml of the fresh culture from each flask were transferred to a new flask. This process was repeated 10 times for 1400 (LNH-ST) and 5 times for 1400 (LNH32) and 1400 (LNH33). The 1400 (LNH-ST) cultures that were cultured for 4, 20, and 40 generations in non-selective medium (each transfer being considered as four generations cultured in non-selective medium) were used to co-ferment glucose and xylose under

- 37 -

similar conditions described in Example 7. The fermentation samples were taken and analyzed identically as described in Example 7. Similarly, the 1400 (LNH32) and 1400 (LNH33) cultures that were cultured for 4 and 20 generations in non-selective medium were used to co-ferment glucose and xylose. Samples were again taken at proper intervals after fermentation was initiated for analysis by HPLC and compared in Figures 4 to 6. These results clearly demonstrate that 1400 (LNH-ST) is far more stable than 1400 (LNH32) and 1400 (LNH33) in maintaining its xylose fermenting capability after being cultured in non-selective medium for more than 40 generations.

EXAMPLE 9

HPLC ANALYSIS OF FERMENTATION SAMPLES

The samples containing the fermentation broth (0.6 ml to 1.0 ml) removed from the cultures were kept in 1.5 ml Eppendorf tubes. The cells and other residues were removed by centrifugation in microfuge (topspeed) for 10 min. The supernatant was diluted 10 fold. The resulting diluted samples were analyzed for its ethanol, glucose, xylose, xylitol, and glycerol contents by high performance liquid chromatography (HPLC), using a Hitachi system according to the following conditions.

- Column: BioRad HPX-87H
- Mobile Phase: 0.005 M H₂SO₄
- Flow Rate: 0.8 ml/min.
- Detection: RI detector
- Temperature: 60°C

- 38 -

• Injection Volume: 20 µl

EXAMPLE 10

5 GENETIC CHARACTERIZATION OF CHROMOSOMAL DNA FROM THE STABLE TRANSFORMANTS 1400 (LNH-ST)

Based on the restriction and PCR analysis, the genetic map (the order and orientation) of the cloned
10 genes, KK, AR, KD, and 5S rDNA present in pLNH-ST, have been determined as shown in Figure 9B. Experiments have been designed to determine whether these genes (KK, AR, and KD) have been integrated into the loci of the 5S rDNA. If these genes have been integrated into the yeast
15 chromosome at the loci of the 5S rDNA as anticipated, the correct size of DNA fragments containing the following combination of partial or intact genes such as 5S rDNA-KK; 5S rDNA-KD; KK-AR, and AR-KD should have been obtained by using 1400 (LNH-ST) chromosomal DNA as the
20 template and the oligonucleotides indicated on the genetic map (Figure 9B) as the primers to carry out DNA synthesis by PCR. If these genes have not been integrated into the yeast chromosome, no such combination of genes or gene fragments should have been obtained by
25 the above described experiments. If these genes have been integrated elsewhere in the yeast chromosome rather than at the loci of 5S rDNA, some of the above described combination of genes or gene fragments should be obtained from the above described experiments, but not those
30 containing the 5S rDNA fragment; such as 5S rDNA-KK and 5S rDNA-KD. For carrying out the above described experiments, chromosomal DNA was isolated from 1400 (LNH-ST), using the protocol provided by Qiagen, Chatsworth,

- 39 -

CA. Positive results were obtained from PCR synthesis by using the following pairs of primers (see Figure 9):

Oligo 25 and Oligo 369; Oligo 26 and Oligo 369; Oligo 370 and Oligo 96; Oligo 97 and Oligo 99; Oligo 982 and

- 5 Oligo27. Thus, based on these analyses, the DNA fragment containing KK-AR-KD seems indeed being integrated in the 1400 yeast chromosome at its 5S rDNA loci.

- 40 -

References

The following publications are indicative of the level of skill possessed by those in the art and are each hereby incorporated by reference as if individually incorporated by reference and fully set forth.

1. Chan, C. S. M., and B.-K. Tye (1980), "Autonomously replicating sequences in *Saccharomyces cerevisiae*," *Proc. Natl. Acad. Sc.*, 77(11), 6329-6333.
2. Cregg, J. M., K. J. Barringer, A. Y. Hessler, and K. R. Madden (1985), "*Pichia pastoris* as a Host System for Transformations," *Molecular and Cellular Biology*, 5, 3376-3385.
3. D'Amore, T., G. Celotto, I. Russell, and G. G. Stewart (1989), "Selection and Optimization of Yeast Suitable for Ethanol Production at 40°" *Enz. Microbial. Technol.* 11, 411-416.
4. D'Amore, T., J. P. Chandra, I. Russell, and G. G. Stewart (1990), "A Study of ethanol Tolerance in Yeast" *Critical Reviews in Biotechnology* 9, 287-304.
5. Ho, N. W. Y., and Tsao, G. T., "Recombinant Yeasts capable of Effective Fermentation of both glucose and Xylose," International Publication No. 95/13362, May 18, 1995, publishing International Application No. PCT/US94/12861, filed November 8, 1994.

- 41 -

6. Kudla, B., and A. Nicolas (1992), "A multisite integrative cassette for the yeast *Saccharomyces cerevisiae*," *Gene* . 119, 49-56.
- 5 7. Kurtz, M. B., M. W. Cortelyou, and D. R. Kirsch (1986), "Integrative Transformation of *Candida albicans*, Using a Cloned *Candida ADE2* Gene," *Molecular and Cellular Biology*, 142-149.
- 10 8. Lopes, T. S., J. Klotwijk, A. E. Veenstra, P. C. van der Aar, H. van Heerikhuizen, H. A. Raué, and R. J. Planta (1989), "High-copy-number integration into the ribosomal DNA of *Saccharomyces cerevisiae*: a new vector for high-level expression," *Elsevier Science Publishers*, 79, 199-206.
- 15 9. Lopes, T. S., Gert-Jan A.J. Hakkaart, B. L. Koerts, H. A. Raué, and R. J. Planta (1991), "Mechanism of high-copy-number integration of pMIRY-type vectors into the ribosomal DNA of *Saccharomyces cerevisiae*," *Elsevier Science Publishers*, 105, 83-90.
- 20 10. Orr-Weaver, T. L., J. W. Szostak, and R. J. Rothstein (1981), "Yeast transformation: A model system for the study of recombination," *Proc. Natl. Acad. Sci.*, 78(10), 6354-6358.
- 25 11. Orr-Weaver, T. L., and J. W. Szostak (1983), "Multiple, Tandem Plasmid Integration in *Saccharomyces*
- 30

- 42 -

cerevisiae," *Molecular and Cellular Biology*, 3(4), 747-749.

12. Romanos, M. A., C. A. Scorer, and J. J. Clare
5 (1992), "Foreign Gene Expression in Yeast: a Review,"
John Wiley & Sons Ltd., 8, 423-488.

13. Rossolini, M., M. L. Riccio, E. Gallo, and C. L.
10 Galeotti (1992), "*Saccharomyces lactis* rDNA as a target
for multiple integration by homologous recombination,"
Elsevier Science Publishers, 75-81.

14. Rothstein, R. J. (1981), "One-Step Gene Disruption
15 in Yeast," *Methods in Enzymology*, 101, 202-211.

15. Rothstein, R. J. (1991), "Targeting, Disruption,
Replacement, and Allele Rescue: Integrative DNA
Transformation in Yeast," *Methods in Enzymology*, 194,
281-301.

20

16. Sakai, A., Y. Shimizu, and F. Hishinuma (1990),
"Integration of heterologous genes into the chromosome of
Saccharomyces cerevisiae using a delta sequence of yeast
25 retrotransposon Ty," *Appl. Microbiol. Biotechnol.*, 33,
302-306.

17. Sakai, A., F. Ozawa, T. Higashizaki, Y. Shimizu, and
F. Hishinuma (1991), "Enhanced Secretion of Human Nerve
30 Growth Factor from *Saccharomyces cerevisiae* Using an
Advanced d-Integration System," *Bio/Technology*, 9, 1382-
1385.

- 43 -

18. Stinchcomb, D. T., M. Thomas, J. Kelly, E. Selker,
and R. W. Davis (1980), "Eukaryotic DNA segments capable
of autonomous replication in yeast," *Proc. Natl. Acad.*
5 *Sci.*, 77(8), 4559-4563.
19. Struhl, K., D. T. Stinchcomb, S. Scherer, and R. W.
Davis (1979), "High-frequency transformation of yeast:
Autonomous replication of hybrid DNA molecules," *Proc.*
10 *Natl. Acad. Sci.*, 76(3), 1035-1039.
20. Szostak, J. W., and R. Wu (1979), "Insertion of a
Genetic Marker into the Ribosomal DNA of Yeast," *Plasmid*,
2, 536-554.
- 15 21. Valenzuela, P., G. I. Bell, A. Venegas, E. T.
Sewell, F. R. Masiarz, L. J. DeGennaro, G. Weinberg, and
W. J. Rutter (1977), "Ribosomal RNA genes of
Saccharomyces cerevisiae II. Physical map and nucleotide
20 sequence of the 5 S ribosomal RNA gene and adjacent
intergenic Regions," *The Journal of Biological Chemistry*,
252(22), 8126-8135.
- 25 22. Chevaller, M. R. and M. Algle (1979), "Qualitative
detection of penicillinase produced by yeast strains
carrying chimeric yeast - Coli plasmid", *Febs Letters*,
108. 179-180.
- 30 23. Ho, N. W. Y., H. C. Gao, J. J. Huang, P. E. Stevis,
S. F. Chang (1984), "Development of a cloning system for

- 44 -

Candida species" *Biotechnol. Bioengineering Symp.* No. 14, 295-301.

24. Ward, A. C. (1990), "Single-steppurification of
5 shuttle vectors from yeast for high frequency back-
transformation into *E. coli*", *Nucleic Acids Research*, 18,
5319.

25. Becker, D. M. and L. Guavente, (1991), "High-
10 efficiency transformation of yeast by electroporation,"
Methods in Enzymology 194, 182-187.

26. Ho, S. N., Hunt, H. D., Horton, R. M., Pullen, J.
K., and Pease, L. R., (1989), "Site-directed mutagenesis
15 by overlap extension using the polymerase chain reaction,
Gene, 77, 51.

27. Kunkel, T. A., (1985), *Proc. Natl. Acad. Sci. USA*,
82, 488.

20 28. Kunkel, T. A., Roberts, J. D. and Zakour, R. A.,
(1987), *Meth. Enzymol.*, 154, 367.

29. Sambrook, J., E. F. Fritsch, and T. Maniatis,
25 (1989), "Molecular Cloning," published by Cold Spring
Harbor Lab. Press, 4.10-4.11.

- 45 -

WHAT IS CLAIMED IS:

1. A yeast which ferments xylose to ethanol, comprising:

5 a yeast having genes integrated at each of multiple reiterated ribosomal DNA sites of the yeast, said genes encoding xylose reductase, xylitol dehydrogenase, and xylulokinase.

10 2. The yeast of claim 1 which also ferments glucose to ethanol.

3. The yeast of claim 2 which is *Saccharomyces*.

15 4. The yeast of claim 3 wherein said sites are non-transcribed DNA sites.

20 5. The yeast of claim 1 wherein the genes are fused to non-glucose-inhibited promoters and the yeast simultaneously ferments glucose and xylose to ethanol.

6. The yeast of claim 5 wherein the promoters do not require xylose for induction.

25 7. The yeast of claim 3 wherein the genes are fused to non-glucose-inhibited promoters and the yeast simultaneously ferments glucose and xylose to ethanol.

30 8. The yeast of claim 4 wherein the genes are fused to non-glucose-inhibited promoters and the yeast simultaneously ferments glucose and xylose to ethanol, the promoters also not requiring xylose for induction.

- 46 -

9. The yeast of claim 6 wherein the xylose reductase and xylitol dehydrogenase genes are from natural yeast which ferment xylose to ethanol.

5

10. The yeast of claim 9 wherein the natural yeast are *Candida Shehatae*, *Pichia stipitis* or *Pachysolen tannophilus*.

10

11. The yeast of claim 9 wherein the xyulokinase gene is from a yeast or bacteria.

12. The yeast of claim 11 wherein the xyulokinase gene is from *Candida Shehatae*, *Pichia stipitis*,

15

Pachysolen tannophilus, *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, or *Escherichia coli*.

13. The yeast of claim 1 having said genes integrated at at least about 10 ribosomal DNA sites of the yeast.

20

14. A method for integrating multiple copies of exogenous DNA into reiterated chromosomal DNA of cells, comprising:

25

(a) transforming the cells with a replicative and integrative plasmid having exogenous DNA including a first selection marker; and

30

(b) repeatedly replicating the cells from step (a) to produce a number of generations of progeny cells while selecting for cells which include the selection marker, so as to promote the retention of the replicative and integrative plasmid in subsequent generations of the

- 47 -

progeny cells and produce progeny cells having multiple integrated copies of the exogenous DNA.

15. The method of claim 14, wherein the plasmid DNA
5 also includes a second selection marker for selecting cells which include the plasmid.

16. The method of claim 14 wherein the cells are
yeast or eukaryotic cells, and wherein the method further
10 includes the step of repeatedly replicating the progeny cells from step (b) to produce a number of generations of progeny cells in the absence of selection for cells which include the selection marker, so as to promote the loss of the plasmid in subsequent generations of progeny cells
15 and recover yeast cells each containing multiple copies of the exogenous DNA integrated into its chromosomal DNA.

17. The method of claim 16 wherein the cells are
yeast cells and the exogenous DNA includes genes encoding
20 xylose reductase, xylitol dehydrogenase, and xylulokinase, which also serve as the first selection marker.

18. The method of claim 14, which comprises:
25 (i) transforming yeast cells with a replicative plasmid having exogenous DNA including a selection marker, the exogenous DNA being flanked on each end by a DNA sequence homologous to a reiterated sequence of DNA of the host;
30 (ii) repeatedly replicating the transformed yeast cells from step (i) to produce a number of generations of progeny cells while selecting for cells which include the

- 48 -

selection marker, so as to promote the retention of the replicative plasmid in subsequent generations of the progeny cells and result in progeny cells each containing multiple integrated copies of the exogenous DNA; and

- 5 (iii) replicating the progeny cells from step (ii) to produce a number of generations of progeny cells in the absence of selection for cells which include the selection marker, so as to promote the loss of the plasmid in subsequent generations of progeny cells and
- 10 recover yeast cells each containing multiple copies of the exogenous DNA integrated into its chromosomal DNA.

19. Yeast cells produced by the method of claim 18.

- 15 20. The yeast cells of claim 19, wherein the exogenous DNA includes genes encoding xylose reductase, xylitol dehydrogenase, and xylulokinase, and the yeast cells ferment xylose to ethanol.

- 20 21. The yeast cells of claim 20, wherein said genes are fused to non-glucose-inhibited promoters which do not require xylose for induction, and wherein the yeast cells ferment glucose and xylose simultaneously to ethanol.

- 25 22. Yeast cells according to claim 21 which substantially maintain their capacity to ferment xylose to ethanol when cultured under non-selective conditions for at least 20 generations.

- 30 23. A yeast which ferments xylose to ethanol, comprising:

- 49 -

a yeast having multiple copies of exogenous DNA integrated into chromosomal DNA of the yeast, the exogenous DNA including genes encoding xylose reductase, xylitol dehydrogenase, and xylulokinase fused to non-glucose-inhibited promoters, the yeast fermenting glucose and xylose simultaneously to ethanol and substantially retaining its capacity for fermenting xylose to ethanol for at least 20 generations when cultured under non-selective conditions.

10

24. The yeast of claim 23, wherein said promoters do not require xylose for induction

15

25. A yeast which ferments xylose to ethanol, comprising:

20

a yeast having multiple copies of an introduced DNA containing genes encoding xylose reductase, xylitol dehydrogenase, and xylulokinase, the yeast fermenting xylose to ethanol and substantially retaining its capacity for fermenting xylose to ethanol when cultured under non-selective conditions for at least 20 generations.

25

26. The yeast of claim 25, wherein the promoters do not require xylose for induction

30

27. A method for fermenting xylose to ethanol, comprising fermenting a xylose-containing medium with a yeast of claim 1, 22, 23, 24, 25 or 26, to produce ethanol.

- 50 -

28. A plasmid vector for integrating an exogenous DNA sequence including a first selection marker into chromosomal DNA of a target yeast cell, the plasmid vector containing a functional yeast DNA replication origin and the exogenous DNA flanked on each end by a DNA flanking sequence which is homologous to a reiterated ribosomal DNA sequence of the target yeast cell, the plasmid further including a second selection marker in a position other than between the DNA flanking sequences.

10

29. A plasmid vector for integrating an exogenous DNA sequence into a yeast to form stable integrants which ferment xylose to ethanol, the plasmid vector containing a functional yeast DNA replication origin and exogenous DNA including genes encoding xylose reductase, xylitol dehydrogenase, and xylulokinase flanked on each end by a DNA flanking sequence which is homologous to a reiterated DNA sequence of the target yeast cell.

15


20

30. A method for forming cells having multiple integrated copies of an exogenous DNA fragment, comprising:

replicating cells having reiterated genomic DNA and which contain a replicative and integrative plasmid containing the exogenous DNA to produce multiple generations of progeny cells while selecting for cells which include the selection marker, so as to promote the retention of the replicative and integrative plasmid in subsequent generations of the progeny cells and produce progeny cells having multiple integrated copies of the exogenous DNA.

25


30



pLNH31, pLNH32, pLNH33, or pLNH 34

Fig. 1


2/12



Fermentation of Glucose and
Xylose by LNH32

Fig. 2


3/12



Simultaneous Fermentation of Glucose and
Xylose by Recombinant *Saccharomyces LNH33*


Fig. 3

4/12




Fermentation of Glucose and Xylose by the
Un-Engineered Parent 1400 *Saccharomyces* Yeast

Fig. 4




Fermentation of glucose and xylose by LNH32
after being cultured for 4 and 20
generarions in non-selective (glucose) medium.

Fig. 5
SUBSTITUTE SHEET (RULE 20)




Fermentation of glucose and xylose by LNH33
after being cultured for 4 and 20
generations in non-selective (glucose) medium.

Fig. 6
SUBSTITUTE SHEET (RULE 26)



Fermentation of glucose and xylose by LNH-ST(1)
after being cultured for 4, 20, and 40
generations in non-selective (glucose) medium.

Fig. 7
SUBSTITUTE SHEET (RULE 20)

Fig. 8

- (A) Yeast (*S. cerevisiae*) AH22 cultured in YEPD (1% yeast extracts, 2% peptone, 2% glucose or YEP (1% yeast extracts, 2% peptone).
- (B) Yeast AH22 cultured in YEPXu (1% yeast extracts, 2% peptone, 2% xylulose) or YEP.

9/12




Fig. 9a





Fig. 9b

SUBSTITUTE SHEET (RULE 26)

10/12

**Fig. 10**


Construction of pKS(-)-KK-AR-KD plasmids

† The XhoI site was regenerated after ligation; *Intact ADC1 promoter;

** ADC1 promoter with TRP5 ribosomal binding site

SUBSTITUTE SHEET (RULE 20)

11/12




Recombinant *Saccharomyces* 1400(LNH-ST)
for fermenting Glucose and Xylose

Fig. 11

SUBSTITUTE SHEET (RULE 26)

12/12

**Fig. 12**

Please type a plus sign (+) inside this box → ☐

PTO/SB/01 (8-96)
 Approved for use through 9/30/98. OMB 0651-0032
 Patent and Trademark Office: U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

DECLARATION FOR UTILITY OR DESIGN PATENT APPLICATION

☒ Declaration OR
Submitted
with Initial Filing

- unsigned

☐ Declaration
Submitted after
Initial Filing

Attorney Docket Number 7024109PUR48

First Named Inventor HO, Nancy W. Y.

COMPLETE IF KNOWN

Application Number

Filing Date

Group Art Unit

Examiner Name

As a below named inventor, I hereby declare that:

My residence, post office address, and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

STABLE RECOMBINANT YEASTS FOR FERMENTING XYLOSE TO EHTANOL

(Title of the invention)

the specification of which

☐ is attached hereto
OR

☒ was filed on (MM/DD/YYYY)

05/06/1997

as United States Application Number or PCT International

Application Number PCT/US97/07663 and was amended on (MM/DD/YYYY) (if applicable).

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment specifically referred to above.

I acknowledge the duty to disclose information which is material to patentability as defined in Title 37 Code of Federal Regulations, §1.56

I hereby claim foreign priority benefits under Title 35 United States Code §119 (a)-(d) or §365(b) of any foreign application(s) for patent or inventor's certificate, or §365 (a) of any PCT international application which designated at least one country other than the United States of America, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or of any PCT international application having a filing date before that of the application on which priority is claimed.

Prior Foreign Application Number(s)	Country	Foreign Filing Date (MM/DD/YYYY)	Priority Not Claimed	Certified Copy Attached?	
				YES	NO
US97/07663	PCT	05/06/1997	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	<input checked="" type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>

☐ Additional foreign application numbers are listed on a supplemental priority sheet attached hereto

I hereby claim the benefit under Title 35, United States Code §119(e) of any United States provisional application(s) listed below

Application Number(s)	Filing Date (MM/DD/YYYY)	<input type="checkbox"/> Additional provisional application numbers are listed on a supplemental priority sheet attached hereto
60/016,865	05/06/1996	

(Page 1 of 5)

Burden Hour Statement: This form is estimated to take 0.4 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Commissioner of Patents and Trademarks, Washington, DC 20231

(January 1997)

Please type a plus sign (+) inside this box → ☐PTO/SB/01 (8-96)
Approved for use through 9/30/98. OMB 0651-0032
Patent and Trademark Office: U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

DECLARATION

I hereby claim the benefit under Title 35, United States Code §120 of any United States application(s), or §365(c) of any PCT international application designating the United States of America, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT International application in the manner provided by the first paragraph of Title 35, United States Code §112, I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations §1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application.

U.S. Parent Application Number	PCT Parent Number	Parent Filing Date (MM/DD/YYYY)	Parent Patent Number (if applicable)
60/016,865	PCT/US97/07663	05/06/1996 05/06/1997	

☐ Additional U.S. or PCT international application numbers are listed on a supplemental priority sheet attached hereto.

As a named inventor, I hereby appoint the following registered practitioner(s) to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith:

Name	Registration Number	Name	Registration Number
Kenneth A. Gandy	#33,386		

☒ Additional registered practitioner(s) named on a supplemental sheet attached hereto.


Direct all correspondence to:

Name	Kenneth A. Gandy at WOODARD, EMHARDT, NAUGHTON, MORIARTY & MCNETT		
Address	Bank One Center/Tower, Suite 3700		
Address	111 Monument Circle		
City	Indianapolis	State	Indiana
Country	US	Telephone	317-634-3456
		Fax	317-637-7561
		ZIP	46204

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true, and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Name of Sole or First Inventor: ☐ A petition has been filed for this unsigned inventor

Given Name	Nancy	Middle Initial	W. Y.	Family Name	Ho	Suffix e.g. Jr.	
------------	-------	----------------	-------	-------------	----	-----------------	--

Inventor's Signature		Date	December 7, 1998
----------------------	---	------	------------------

Residence: City	West Lafayette	State	IN	Country	US	Citizenship	US
-----------------	----------------	-------	----	---------	----	-------------	----

Post Office Address	606 Riley Lane		
---------------------	----------------	--	--

Post Office Address	West Lafayette, Indiana 47906		
---------------------	-------------------------------	--	--

City	West Lafayette	State	IN	Zip	47906	Country	US
------	----------------	-------	----	-----	-------	---------	----

☒ Additional inventors are being named on supplemental sheet(s) attached hereto

Please type a plus sign (+) inside this box → ☐

PTO/SB/01 (8-96)
 Approved for use through 9/30/98. OMB 0651-0032
 Patent and Trademark Office: U.S. DEPARTMENT OF COMMERCE
 Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

+

DECLARATION					ADDITIONAL INVENTOR(S) Supplemental Sheet				
Name of Additional Joint Inventor, if any:					<input type="checkbox"/> A petition has been filed for this unsigned inventor				
Given Name	<u>Zheng-Dao</u>	Middle Initial		Family Name	<u>Chen</u>	Suffix e.g., Jr.			
Inventor's Signature	<u>Zhengdao Chen</u>				Date	<u>December 7, 1998</u>			
Residence: City	<u>West Lafayette</u>	State	<u>IN</u>	Country	<u>US</u>	Citizenship	<u>CN</u>		
Post Office Address	<u>480 Maple Street</u>								
Post Office Address	<u>West Lafayette, Indiana 47906</u>								
City	<u>West Lafayette</u>	State	<u>IN</u>	Zip	<u>47906</u>	Country	<u>US</u>		
Name of Additional Joint Inventor, if any:					<input type="checkbox"/> A petition has been filed for this unsigned inventor				
Given Name		Middle Initial		Family Name		Suffix e.g., Jr.			
Inventor's Signature					Date				
Residence: City		State		Country		Citizenship			
Post Office Address									
Post Office Address									
City		State		Zip		Country			
Name of Additional Joint Inventor, if any:					<input type="checkbox"/> A petition has been filed for this unsigned inventor				
Given Name		Middle Initial		Family Name		Suffix e.g., Jr.			
Inventor's Signature					Date				
Residence: City		State		Country		Citizenship			
Post Office Address									
Post Office Address									
City		State		Zip		Country			
Name of Additional Joint Inventor, if any:					<input type="checkbox"/> A petition has been filed for this unsigned inventor				
Given Name		Middle Initial		Family Name		Suffix e.g., Jr.			
Inventor's Signature					Date				
Residence: City		State		Country		Citizenship			
Post Office Address									
Post Office Address									
City		State		Zip		Country			

☐ Additional inventors are being named on supplemental sheet(s) attached hereto

Please type a plus sign (+) inside this box →

PTO/SB/01 (8-96)


Approved for use through 9/30/98. OMB 0651-0032
Patent and Trademark Office: U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

DECLARATION	PRIORITY DATA (Supplemental Sheet)
--------------------	--

Additional foreign applications:					
Prior Foreign Application Number(s)	Country	Foreign Filing Date (MM/DD/YYYY)	Priority Not Claimed	Certified Copy Attached?	
				YES	NO
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Additional provisional applications:					
Application Number		Filing Date (MM/DD/YYYY)			
Additional U.S. applications:					
U.S. Parent Application Number	PCT Parent Number	Parent Filing Date (MM/DD/YYYY)	Parent Patent Number (if applicable)		

United States Patent & Trademark Office
Office of Initial Patent Examination -- Scanning Division



Application deficiencies were found during scanning:

☒ Page(s) 5 of Declaration were not present
for scanning. (Document title)

☐ Page(s) _____ of _____ were not present
for scanning. (Document title)

☐ Scanned copy is best available.

SCANNED5